


REVIEW ARTICLE

A review on the Immunological Techniques Use for Detection of *Schistosoma* spp Infection

¹Suleiman, Jafaru ; ²Isyaku, Nike Tawakaltu; ²Ukatu, Victoria E.; ¹Yusuf, Ibrahim Alhaji; and ³Abubakar, Hafiz

¹Department of Biological Sciences, Faculty of Science, Sokoto State University, Sokoto State, Nigeria

²Department of Animal and Environmental Biology, Faculty of Life Sciences, Kebbi State University of Science and Technology Aleiro, Nigeria

³Department of Biological Sciences, Faculty of Science, Federal University Birnin Kebbi, Kebbi State, Nigeria

ABSTRACT

Over 200 million people are suffering from the debilitating *Schistosomiasis*, which has the greatest morbidity and mortality rates in African nations. *Schistosomiasis* continues to be a tropical disease that receives little attention from governments and healthcare Institutions while having a wide range of negative impacts on society's health and socioeconomic hardship. The establishment of precise investigation for both gut and urinary *Schistosomiasis* is one of the crucial areas that are severely undeveloped. A review was conducted to highlight immunological methods used to detect *Schistosomiasis*. Databases from Science Direct, World Health Organization, and PubMed were used. Articles for which at least the abstract was available in English were selected for the present study. Relevant articles were screened, duplicates were eliminated, eligibility standards were followed, and qualified studies were reviewed. Techniques including Skin Reaction Test/Biopsy, Indirect Immuno-Fluorescence Test (IFT), Indirect Hemagglutination Test (IHAT), Circumoval Precipitin Test (Copt), Monoclonal Antibodies Test (Mabt), ELISA, and others were discussed. The challenges faced by these techniques were also highlighted. Among the techniques discussed, ELISA was found to be the most widely used as it is the most effective, easy to use, and could detect both chronic and acute *Schistosomiasis*

ARTICLE HISTORY

Received: September 12, 2022

Accepted: September 27, 2022

Published: September 30, 2022

KEYWORDS

Immunology, Diagnosis, Schistosomes, Infection



© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

INTRODUCTION

The physiological and socioeconomic well-being of individuals in vulnerable communities is severely harmed by schistosomiasis, with school-aged children and adults being the most infected groups with the highest morbidity and mortality rate (Infurnari *et al.*, 2017). Six species, namely *Schistosoma mansoni*, *S. haematobium*, *S. japonicum*, *S. mekongi*, *S. guineensis*, and *S. intercalatum*, are known to cause *schistosomiasis* globally (Chernet *et al.*, 2017). In Sub-Saharan Africa, *S. mansoni* (agent of intestinal schistosomiasis) and *S. haematobium* (agent of urogenital schistosomiasis) are the two main schistosome pathogens (Cioli and Pica-Mattoccia, 2013). Host reactions to schistosome eggs cause the chronic disease's clinical symptoms (Gryseels *et al.*, 2016). Eggs from *S. mansoni* and *S. japonicum* most frequently become lodged in the liver or intestine's blood vessels, resulting in diarrhea, constipation, and blood in the urine (John, 2014). In cases of severe infections, liver fibrosis and portal hypertension may also occur as a result of systemic inflammation,

which can also cause gut wall ulceration, hyperplasia, and peptic ulcer disease (Lindholz *et al.*, 2018).

Infections with *Schistosoma* species are mostly identified through microscopic analysis of the parasites' eggs in urine or stool samples taken from infected individuals (Akinwale *et al.*, 2011). However, the immunological method develops are more quickly, and technically cheaper, than microscopic examinations (Cioli and Pica-Mattoccia, 2013). Aside from providing tests with high sensitivity and specificity, the immunological approach in epidemiologic studies also evaluates the efficacy of various control methods in regions where parasitic infections are prevalent (Barakat, 2013). These tests can be employed even when parasites are reproducing (Hamburger *et al.*, 2017).

Since some schistosome-infected individuals may exhibit clinical symptoms in endemic areas, a review of the immunological diagnosis is crucial for the present situation. The methods also lead to the development of

Correspondence: Suleiman J. Department of Biological Sciences, Faculty of Science, Sokoto State University, Sokoto

✉ Suleiman.jafar@ssu.edu.ng

How to cite: Suleiman J., Isyaku N. T. Ukatu, V. E., Alhaji, Y. I. and Abubakar, H. (2022). A review on the Immunological Techniques Use for Detection of *Schistosoma* spp Infection. *UMYU Scientifica*, 1(1), <https://doi.org/10.56919/usci.1122.039>

tools that allow treatment monitoring and also assessing the effectiveness of chemotherapy as well as other curative interventions (Hervé *et al.*, 2013; Devaney *et al.*, 2010). Rates of Immunoglobulin E (IgE) to the Schistosomes peptides have been associated in some regions of the world with susceptibility to superinfection after treatment (Pearson *et al.*, 2012). The effectiveness of vaccine programs can be anticipated in the future when anti-parasitic vaccination is available (Pennington *et al.*, 2017). The Discovery of this effective vaccine could be greatly influenced by a study of the methods for detecting these antigens since it will aid in identifying those people who develop immunity to parasite infection (Fitzsimmons *et al.*, 2005).

Immunological Diagnosis

The identification of *Schistosomiasis* is mostly based on conventional methods (mostly microscopic examination) (Gray *et al.*, 2021). However, the identification of adult worms and antigens in serum or urine has a lot of diagnostic capabilities and it may eventually replace the conventional techniques (Hervé *et al.*, 2013). The antigens found in serum or urine are either cathodic (circulating cathodic antigen) or anodic (circulating anodic antigen) (Pennington *et al.*, 2017). The most common methods include the skin reaction test/biopsy, indirect immunofluorescence test (IFT), indirect hemagglutination test (IHAT), circumoval precipitin test (COPT), various monoclonal antibody tests (MAbT), and various e-PCR formats (Deelder, 2013; Meurs *et al.*, 2018; Knopp *et al.*, 2019). For the diagnosis of *Schistosomiasis*, some immunological techniques have been put forth, but only a select handful have been able to advance to the large field and confirmation steps (Luciana *et al.*, 2013).

Skin Reaction Test/Biopsy

Skin Reaction Test also known as "Biopsy" comes from the Greek words "bio-life" and "opsia," to see," (Patton *et al.*, 2018). Technically, Biopsy is the removal of tissue from living things for microscopic analysis and diagnosis (Sanjay *et al.*, 2014). Ernest Besnier first used the term "biopsy" in medical terminology in 1879 (Karkera *et al.*, 2011). The primary goal of a biopsy is to confirm a clinical and radiographic diagnosis, guide surgical treatment, and verify whether a lesion has been completely removed (Kumaraswamy *et al.*, 2012). It is also helpful in choosing the type of treatment to implement for specific disorders (Catherine *et al.*, 2008). If necessary, biopsy reports can also serve as medical records (Karkera *et al.*, 2011). In China during the middle of the 1950s, *Schistosoma* adult worm skin test antigen was employed regularly, and various epidemiological studies on skin reaction tests were also carried out during the years 1963, 1969, and 1976 (Smithers and Terry, 2015). However, it was discovered that test results remained low also when given a good prediction (Lalonde *et al.*, 2015).

The method primarily usually used for skin biopsies in the detection of *Schistosoma* infection is a punch biopsy

(McManus *et al.*, 2010). This method can be applied to both therapeutic and diagnostic objectives (viz, punch excision of small pyogenic granuloma, verruca, tattoo) (Guegan *et al.*, 2019). Any hard lesion and microscopic vesicles that fit inside the perforation can be treated using this technique (Thomson *et al.*, 2017).

With a disposable or sterilizable punch of various sizes, the lesion or region to be biopsied is removed. A 4-mm punch is acceptable for non-facial lesions (Gray *et al.*, 2011). However, biopsies of 5 mm or greater are preferred in granulomatous disorders or conditions with unusual features (Friedman, 2004). It is advised against taking biopsies that are less than 3 mm since important features could be overlooked (Krunic *et al.*, 2014).

Just after the region has been put under anesthesia, the skin is pulled in a perpendicular direction to the lines of the skin that is at rest. With this technique, the round skin defect is transformed into an ellipse, reducing the possibility of dogear development after suturing (Rabello *et al.*, 2012). The thumb, middle finger, and index finger are used to support, rotate, and apply pressure to the punch, whereas the index finger stabilizes and applies pressure to the punch (Wilhelmi *et al.*, 2011). Pressing the punch into the skin with rotating motions causes a sensation to be felt (Zhu, 2015). This is due to punching penetration into the gelatinous mass subcutaneous tissue plane (Gray *et al.*, 2011). The wound that results from this can either be sutured or left untreated to allow for further healing (Krunic *et al.*, 2014). The face, genitals, and mucosa, have excellent flow, heal fast, and leave little scars (Perez *et al.*, 2013).

It's crucial to avoid hitting the periosteum when doing a biopsy on the forehead, shin, dorsum of the nose, or scalp because doing so produces excruciating pain and agony (Mutalik, 2018). Punch biopsy has the benefits of being simple to do and collecting evenly shaped tissues, but the drawbacks include the possibility of insufficient material being retrieved and the tendency for the biopsy to exclude deeper tissue (Zhu, 2015).

Indirect Immuno-Fluorescence Test (IFT)

Indirect Immuno-Fluorescence Test is a common method for detecting antibodies, the indirect immunofluorescence antigens (IFA), detects antibodies by their unique ability to bind to parasitic antigens produced in infected cells (Vendrame *et al.*, 2021). IFT is also called the Secondary immunofluorescence test used as the laboratory technique for identifying circulating autoantibodies in patient serum also a method is employed to identify autoimmune disorders that rupture (Grenfell *et al.*, 2013). In this form of test bound antibodies are visible after being incubated with fluorescently labeled antihuman antibodies (Zhu, 2015). IFT is one of the cell imaging methods which rely on the use of antibodies to mark a particular target antigen with a fluorescent dye called fluorescein isothiocyanate (FITC) (Smith *et al.*, 2012).

In IFT, primary conjugated antibodies with chemically labeled fluorophores are frequently utilized (Grenfell *et al.*, 2013). Two antibodies are used in indirect IF; the primary antibody is unconjugated, while the secondary antibody is directed against the primary antibody and conjugated to a fluorophore for detection (Deelder *et al.*, 2020). Studies have shown that an indirect immunofluorescence reaction, using paraffin-embedded sections of adult worms, allows for the detection of immunoglobulin M (IgM) antibodies against antigens of the parasites' digestive tract, making it a highly sensitive method for the diagnosis of acute and chronic schistosomal infections (Vendrame *et al.*, 2021). The specificity of this reaction has also proved to be adequate (Smith *et al.*, 2012).

In 1998, in a low-endemic region of Brazil, immun-epidemiologic studies of *Schistosoma mansoni* infection used IFT as a diagnostic tool (Zhu, 2015). It was discovered that stomach fluorescence, which can appear forty-four (44) days after infection and is always connected to acute *Schistosoma mansoni* infection, can be detected by IFT using adult worm segments (Kanamura *et al.*, 2019). Once fluorescence is found in the parenchyma and/or stomach of the worms, frozen sections of a sample are classified as positive for the igg Antibodies immunofluorescence test (IgG-IFT), and as positive for the IgM-indirect immunofluorescence test (IgM-IFT) (paraffin sections) when fluorescence is found in only the gut of the worms (Burlandy-Soares *et al.*, 2013).

Indirect immunofluorescence has the benefits of high sensitivity and easy signal color change dependent on different second antibodies that may be purchased commercially; it is simple to obtain the second antibodies that have been tagged (Alem *et al.*, 2017). The advantages of indirect immunofluorescence include high sensitivity and simple signal color change depending on various second antibodies that may be bought commercially; the second antibodies that have been labeled are easy to obtain, High sensitivity and a quicker processing time are two advantages indirect immunofluorescence over direct immunofluorescence; direct immunofluorescence must be performed when there are several antibodies from the same species (Eriksson *et al.*, 2011). The use of direct immunofluorescence has some drawbacks; compared to indirect immunofluorescence, the first has a lesser signal; the second is that direct immunofluorescence is more expensive than indirect immunofluorescence because the labeling process is more complicated and less flexible when there are no commercially available direct conjugates (Hargraves, 2019).

The primary drawback of the indirect immunofluorescence method is the inability of the anti-immunoglobulin sandwich reagent to differentiate between exogenous and endogenous immunoglobulin (Lodh *et al.*, 2013). For instance, independent of the presence or absence of additional antibodies, a sandwich reagent of rabbit anti-mouse immunoglobulin would

detect membrane immunoglobulin in mouse B cells. Since rabbit or rat antimouse immunoglobulin will only weakly or completely cross-react with human immunoglobulin, the issue is not as serious when mouse monoclonal antibodies are employed to stain human cells (Aryeetey *et al.*, 2013). By passing over human immunoglobulin-coupled beads, antihuman antibodies from several commercial antisera to mouse immunoglobulins have been removed (Carvalho *et al.*, 2012). Nevertheless, there is a chance for the cross-reactions, hence it is necessary to do controls in which the first antibody is not used (Enk *et al.*, 2012).

Indirect Hemagglutination Test (IHAT)

An indirect hemagglutination test is a form of immunological test in which an antigen is entrapped onto erythrocytes of a specimen and agglutination of the antigen-sensitized erythrocytes is induced by the action of an antibody specific to the adsorbed antigens, said method being characterized by the removal of heterologous agglutinins from the serum to be tested that may react with the homologous red blood cells used (Ahmed, 2019).

The surface proteins that agglutinate red blood cells (RBC) of diverse species are encoded by the nucleic acids of different Schistosomes (Whitty *et al.*, 2018). Hemagglutination is the process by which Schistosoma hemagglutinins interact with red blood cells to produce a lattice of agglutinated cells that settle unevenly in a tube or microtiter well and unagglutinated cells condense into a button-like structure (Erkol *et al.*, 2009).

The procedure for the hemagglutination test involved; obtaining a parasites preparation with a known HA titer, of the particular parasite, or determining the HA titer of the parasites, then preparing two-fold dilutions of the patient/test serum, say from 1:4 to 1:1024 for the test, except for the serum control wells, add a constant quantity of parasites to each well on a 96-well plate that is equal to 4 HA units, 60 minutes should be given for the plate to stand at room temperature (time varies according to specific requirements); Red blood cells (RBC) are added, and 30 minutes are spent incubating at 4 °C then check out the wells (Ortu *et al.*, 2017).

The IHAT kit's reagents and samples must be allowed to reach room temperature before the test is conducted (serum will be separated from a blood sample by centrifugation) (Azab and El Zayat, 2016). A disposable tube containing 0.05 ml of test serum and 1.95 ml of buffer solution must be used to deliver the 1/40 stock dilution of test serum (El Ganayni *et al.*, 2012). Using a micro pipettor, deliver 50µL of buffer solution into the wells. After setting up the microplate widthwise, add 50µL of serum stock dilution in the first well that was mixed with buffer. Next, transfer the 2µL from the first well into the second well, preferably using a microdilution. Finally, discard 50µL into the wells before adding 50 µL of the

serum stock-dilution (from step 1) and mixing it with buffer, the 1:80 dilutions is used as "serum control," which helps to prevent the occurrence of natural anti-sheep agglutinins in some sera (Feldmeier *et al.*, 2013).

Suspensions of red blood cells gently jiggle One drop of non-sensitized red blood cells was added to the seventh well (serum control), one drop of sensitized red blood cells was added to the eighth well (reagent control), and one drop of sensitized red blood cells was added to the ninth well. One "reagent control" was set up per microplate, and the content of the well was very carefully homogenized by lateral thrumming on the edge (do not use orbital vibration). Read the response after two hours (El Ridi *et al.*, 2016).

The Indirect Haemagglutination (IHA) strategy utilizes different extracts of red blood cells encapsulated with an enzyme, typically straightforward solubilized egg antigen (SEA), for the identification of positive patient sera via agglutination (Gui *et al.*, 2019). Although this method is not currently used, it was used in Chinese population studies as a substitute test to identify human infection with *S. haematobium*. Its social acceptability is also limited, and because of its poor or variable precision, which renders it inappropriate for personal management and therapy, it can only be advised as an additional technique for or before usages (Yu *et al.*, 2017).

Few researchers have previously studied indirect haemagglutination (IHA) tests employing adult worm antigens for *Schistosomiasis* diagnosis; Most of the previously utilized IHA test kits were not commercially accessible for use during Epidemiological application, with a few exceptions, Sensitivity and specificity were reported in these earlier research to range from 71 to 100% and 80 to 100%, respectively (Van Gool *et al.* 2002).

The identical commercial kit IHAT developed by Fumouze Laboratories was evaluated by (Tom-Van *et al.*, 2002) for the identification of *Schistosoma haematobium*; the study's sensitivity was 80% and its specificity was 98.9%. The study used a cutoff of 1/160 and included only 25 *S. haematobium*-infected cases (Van Gool *et al.*, 2002); Kinkel *et al.* (2012) discovered that the IHA's sensitivity for *S. haematobium* was 71.4% and specificity was 99.0%.

Some researchers used a commercial IHA kit (Cellognost *Schistosomiasis* H) manufactured by Dade Behring Marburg GmbH, Germany, and the test was carried out according to the manufacturer's instructions, it was reported that the sensitivity of the IHA test for the diagnosis of *Schistosomiasis mansoni* was found to be 83%, while the specificity was lower (53%) (Berhanu Erko *et al.*, 2009).

Circum Oval Precipitin Test (COPT)

One of the primary techniques for use in the immunodiagnosis of *Schistosomiasis* is the Circumoval Precipitin (COP) test, which was first described by Oliver Gonzalez (1954). (Kagan and Pellegrino, 2016). Although

the method is very straightforward, it was further simplified by the use of lyophilized eggs (Revera de Sala *et al.*, 2012; Yogore *et al.*, 2018). These updates were thought to be quite useful and appropriate for surveys in rural areas (Nosenas *et al.*, 2015; Tanaka *et al.*, 2015; Matsuda *et al.*, 2017).

The paddock COP test has also become more useful as a result of the recent research on the availability of air-dried *S. japonicum* eggs (Kamiya, 2013). Additionally, a noteworthy article on the use of microscope slides with perforated Dubl-style tape describes how lyophilized eggs were exposed to test blood and how the reaction was viewed via the microscope coverslips (Lewert and Yogore, 2019).

For the COP test, a roll of PVC electrical insulating tape is drilled with heated 11 mm and 5 mm tubular-cork borers, the two holes are positioned so that they have a common edge that opens to each other (Tanaka *et al.*, 2015). The slide is normally left in a room for at least 2 days to ensure tight adhesion of the tape; otherwise, leakage would occur through minute gaps. Then, about 10 l of *S. japonicum* egg suspension containing roughly 100 eggs obtained from an 8-week infected rabbit is placed in the large hole using a micropipette, spread evenly, and allowed to dry (Yokogawa *et al.*, 2016); this procedure is described by Kamiya (2013), A drop of serum will be placed in the large hole at the edge enclosing the small hole, and a piece of cellulose tape (24 mm wide, 35 m long, 0.02 mm thick, Sekisui Co. Ltd., Japan) is applied to the insulating tape starting from the side of the large hole toward the small hole with continuous gentle pressing with the index and middle fingers, this procedure caused the serum to enter the space between the tape and the slide (Tanaka *et al.*, 2015).

According to preliminary investigations, just a single hole would not be sufficient because the adhesiveness of the tape is diminished when too much serum is applied (Nosenas *et al.*, 2015). Therefore, a smaller hole is required as a reservoir for any overflow (Lewert and Yogore, 2019). After the final slide has been incubated for 48 hours at room temperature, the COP response is microscopic and examined using cellulose tape (Nosenas *et al.*, 2015).

The circumoval precipitin test (COPT) is an immunological test based on the precipitation of patient serum with lyophilized eggs or purified live eggs identified under a microscope, is helpful for the prognosis of *S. mansoni* and *S. japonicum* due to its high sensitivity (92-100%) and specificity (96-100%) (Zhou *et al.*, 2017); this method was used in China and Venezuela to identify low transmission zones, together with other coproscopic and serologic testing, in a modified form as a component of a commercial product (Dunne *et al.*, 2014). The disadvantages of this test include that it takes a very long time (48 hours), is difficult to perform, and has a variable period of seroconversion after treatment, according to the

experiences reported for Venezuela (negative rate of 64% of patients after 12 months of praziquantel treatment) and China (negative rate of 0% after 12 months) (Alarcón *et al.*, 218).

Serologic assays have shown clinical value for diagnosis by the identification of antibodies against schistosomal antigens due to the exceptionally broad spectrum of the reported immunodiagnostic assays (Doenhoff *et al.*, 2013); this approach is particularly useful for symptomatic travelers or serological surveys (Webster *et al.*, 2019). Current serologic tests cannot differentiate between active infection and prior exposure in people residing in *Schistosomiasis*-endemic locales, although several isotypic assays can frequently group active or inactive illnesses (Dunne *et al.*, 2014).

Since many years ago, monoclonal antibodies have been employed to identify circulating schistosomal antigens, which has the advantage of semi-quantitatively identifying active infections (Hagan *et al.*, 2011). A point of contact circulating cathodic antigen (POC-CCA) assay is now available from Rapid Medical Diagnostics, Pretoria, and RSA to map *Schistosoma* infections (Colley *et al.*, 2017). This parallel flow cassette test allows for on-site tracking of *S. mansoni* without stool samples, and it looks to be more sensitive than the Kato-Katz assay for mapping *Schistosoma* endemic areas (Rihet *et al.*, 2017). This assay will be an essential tool for conducting control programs in new sites (Towbin *et al.*, 2019).

Monoclonal Antibodies Test (MABT)

Using specific monoclonal antibodies (MAbs) to identify circulating schistosome antigens secreted by living schistosomes in bodily fluids has while back been shown to be a hopeful technique for detecting active infection, assessing the effectiveness of treatment, and determining the efficacy of upcoming vaccines (Van Etten *et al.*, 2014). The antigen detection assays' typically high levels of sensitivity have been confirmed by contrasting their findings with those of quantitative parasitological techniques (Deelder *et al.*, 2014).

Subjects demonstrated a sensitivity of 80 to 90% when excreting at least 100 eggs per gram (EPG) of feces, and 100% when excreting at least 400 EPG. Techniques for detecting antigens, all of which rely on the use of MAbs, have nearly 100% specificity (Polman *et al.*, 2015). Numerous antigen error-checking assays have both high specificities and high sensitivities; however, because they need costly, special tools and bureaucratic procedures, it is challenging to adapt them for use in the field (Nash, 2018).

Enzyme-Linked Immunosorbent Assays (ELISA)

It is mentioned that one form of immunodiagnostic test that can be used to find *Schistosomiasis* is the dot enzyme-linked immunosorbent assay (ELISA). To establish a more field-applicable test format, many changes have been recorded (Ambrose *et al.*, 2018). The most used test

for *Schistosomiasis* diagnosis is the ELISA, which allows for detecting many antibody classes and uses several antigens (Stete *et al.*, 2019).

Low specificity was discovered in early investigations on the detection of *Schistosoma* antigens, which prompted researchers to look for pure antigen preparations such as cationic fraction 6 (CEF6), adult microsomal antigens for *S. mansoni* (MAMA), *S. japonicum* (JAMA), and *S. haematobium*. Hemocyanin from keyhole limpets, gut-associated antigen (KLH), crude soluble egg antigens (SEA), and soluble adult (HAMA) were initially used for the detection of *Schistosoma* antigens (Olliaro *et al.*, 2020).

CEF6 is an isolated antigen from *Schistosoma* eggs that, compared to other antigens, has the highest association with egg production and an ELISA platform that demonstrated 91.7% sensitivity and 90% specificity (Dunne *et al.*, 2017). Three field investigations conducted in *S. mansoni* endemic regions of Kenya (Doenhoff *et al.*, 2016), Burkina Faso (Sorgho *et al.*, 2015), and Saudi Arabia since the publication of that article have continually assessed this antigen (Ghandour *et al.*, 2017).

The accuracy was still low, though (59%, 55%, and 17.2%, respectively). All of these investigations raised concerns about the usefulness of the reference tests, and the low specificity was attributed to the Kato-Katz method's poor sensitivity, which served as the gold standard. All three assessments' sensitivity levels were confirmed to be high (97%, 90%, and 97%, respectively), but their specificity was still low (59%, 55%, and 17.2%) (Luciana *et al.*, 2013).

According to a study conducted in Egypt employing MAMA and HAMA as screening tests in a region where *S. haematobium* has been endemic in which MAMA FAST-ELISA had a responsiveness of 83.6% and HAMA FAST-ELISA had an attentiveness of 98.5%. All egg-passing subjects passed the test using the HAMA FAST-ELISA together with the Western blot method for additional examination. Combining adult worm mitochondrial antigens with both the Falcon assay screening test enzyme was explored (Al-Sherbiny *et al.*, 2019).

A systematic serosurvey involving nearly all Puerto Rican Towns investigated 2,955 normal healthy blood samples using MAMA FAST ELISA and the results showed that 15.4% of samples were positive and 10.6% had EITB confirmation, the highest seroprevalence rates (21.1-38.5%) were concentrated in 17 municipalities which together showed 48.8% of all seropositive samples and 18.8% of all samples with EITB confirmation (Tsang *et al.*, 2021).

The FAST-HAMA test demonstrated a positive rate of 95% in recent research, showing that it can be used as an application required for *S. haematobium* contamination. This test uses a purified HAMA antigen that is highly

specific for *S. haematobium* and a unique standard curve. Additional research is required to confirm the assay's specificity and usefulness in real-world settings (Abdel-Fattah *et al.*, 2018).

Epidemiological data on *S. mansoni* in Sudan and *S. japonicum* in China found high sensitivity but low specificity for kilodalton adult *Schistosoma* antigens; this low specificity was justified by some factors, including cross-reaction, infection with non-human *Schistosoma* species, and lack of sensitivity of the parasitological reference test; regrettably, additional assessments of the accuracy of this methodology are not available, and this issue continues (Wen *et al.*, 2019).

Other Immunological Methods

Quick tests for the detection of antibodies for *S. japonicum* include the dot immunogold filtration assay, the silver-enhanced colloidal gold metalloimmuno assay, and the colloidal dye immune filtration assay, similarly, an immunoblot test of the *S. mansoni*, *S. intercalatum*, and *S. haematobium* membrane antigen has also been examined for the detection of antibodies against *S. mansoni* (Cesari *et al.*, 2018).

Advantages and Drawbacks of Immunological diagnosis

Numerous different immunological methods have some common shortcomings, including sensitivity (depending on the standard used), intrinsic test characteristics (such as antigen and erythrocyte type), the prevalence of infection in the investigated community, the potential for positive reactions to persist after treatment, and cross-reactions with other helminths (Sorgho *et al.*, 2015).

Although immunological detection methods are quite effective, they are not frequently utilized for epidemiological research and do not form part of the diagnostic process for *Schistosomiasis* (Cesari *et al.*, 2018). Immunological methods are frequently criticized for their lack of specificity and inability to discriminate between an infection that is present today and one that was present in the past (Utzinger *et al.*, 2015; Worrell *et al.*, 2015). Alternative hypotheses for the lack of specificity, however, include the possibility that egg-negative, antibody-positive patients may have infections that are missed by imprecise parasitological techniques that failed to detect eggs in subjects with mild infections (Coulibaly *et al.*, 2013; Hawkins *et al.*, 2014), or who received sub-

curative drug doses (King, and Bertsch, 2014; Hinz *et al.*, 2017). Of course, there is also a chance that false-positive antibody tests are brought on by antigens from other schistosomes (Abdel-Fattah *et al.*, 2018). Another drawback is the gradual drop in particular antibody levels following therapy (Corstjens *et al.*, 2014). Additionally, in endemic locations, blood collection is difficult to implement under field conditions (Van Lieshout *et al.*, 2017). Despite these drawbacks, antibody detection is still likely to be the best diagnostic tool for schistosome infection in low-infected areas for some time to come. Antibody tests may also be useful for monitoring *Schistosomiasis* exposure in places that are moving toward elimination (Barda *et al.*, 2013).

Moving Forward

Most developing nations still rely mainly on microscopy for the diagnosis of intestinal and urogenital *Schistosomiasis*. Although this is the WHO-approved method for treating urogenital *Schistosomiasis*, many cases of intestinal *Schistosomiasis* as well as urogenital schistosomiasis go unreported in nations with low rates of schistosome transmission. For stakeholders and partners, finding affordable diagnostics for highly sensitive conditions is a difficult obstacle to overcome. Africa's endemic nations will not be able to afford monitoring kits without assistance from the international community. As we move closer to the suggested eradication of Schistosomes, this poses a barrier to tracking the effectiveness of control measures being implemented. It is important to put more effort into creating RDTs that are less expensive, more sensitive, and more targeted, especially for *S. haematobium*. Last but not least, a lower price review is required for the ELISA test kit, to make it more widely applicable and useful in endemic areas.

CONCLUSION

Immunological detection is the form of diagnosis that involved the use of reagents experimental approach as diagnostic tools and approach to research the immune system, the most popular techniques used for the detection of *Schistosomiasis* infection are Biopsy, IFT, IHAT, COPT, MAbT, and ELSA; Although immunological methods are more specific and sensitive than conventional methods ELISA was more often used while conducting immune epidemiological studies of *Schistosomiasis*.

Ahmed, A. Y. (2019). Validity of indirect haemagglutination assay (IHA) in the detection of *Schistosoma haematobium* infection relative to microscopic examination. *Journal of Medical and Life Science*, 1(2); 57 -64. [Crossref]

Akinwale, O. P., Kane, R. A., Rollinson, D., Stothard, J. R.; Ajayi, M. B., Akande, D. O., Ogungbemi, M. O., Duker, C., Gyang, P. V., and Adeleke, M. A.

REFERENCES

Abdel-Fattah, M., Al-Sherbiny, M., Osman, A., Charmy, R., and Tsang, V. (2018). Improving the detection limit of quantitative diagnosis of anti-*S. haematobium* antibodies using Falcon Assay Screening Test (FAST) ELISA by developing a new standard curve. *Journal of Parasitological*

- (2011). Molecular approaches to the identification of *Bulinus* species in south-west Nigeria and observations on natural snail infections with schistosomes. *Journal of Helminthology*, 85, 283-293. [[Crossref](#)]
- Alarcón de N. B., Ruiz, R., Losada, S., Colmenares, C., Contreras, R., and Cesari, I. M. (2018). Detection of Schistosomiasis cases in low-transmission areas based on coprologic and serologic criteria The Venezuelan experience. *Acta Tropica*; 103:41-49. [[Crossref](#)]
- Alem, M., Moghadam, S., Maliki, J., Zaidi, A., Nayak, N., and Li, T. M. (2017). Detection of autoantibodies to nuclear antigen by EIA and IF techniques. *Journal of Allergic Immunology*, 188: 191-194.
- Al-Sherbiny, M. M., Osman, A. M., Hancock, K., Deelder, A. M., and Tsang, V. C. (2019). Application of immunodiagnostic assays: detection of antibodies and circulating antigens in human Schistosomiasis and correlation with clinical findings. *American Journal of Tropical Medicine and Hygiene*; 60:960-966. [[Crossref](#)]
- Ambroise, T. P., Loizzo, T., and Desgeorges, P. T. (2018). Human Schistosomiasis due to *Schistosoma mansoni*, *S. haematobium* and *S. japonicum*. Serological diagnosis by ELISA, immunofluorescence, and indirect hemagglutination]. *Annal of Sociology in Belgium Medical Tropic.*; 61: 379-392.
- Aryeetey, Y. A., Essien-Baidoo, S., Larbi, I. A., Ahmed, K., Amoah, A. S., Obeng, B. B., (2013) Molecular diagnosis of *Schistosoma* infections in urine samples of school children in Ghana. *American Journal of Tropical Medicine and Hygiene*; 88:1028-1031. [[Crossref](#)]
- Azab, M. E., and el Zayat, E. A. (2016). Evaluation of purified antigens in haemagglutination test (IHA) for determination of cross reactivities in diagnosis of fascioliasis and Schistosomiasis. *Journal of Egyptian and Sociological Parasitology*. 26:677-685.
- Barakat, R. M. (2013). Epidemiology of Schistosomiasis in Egypt: travel through time. *Journal of Advance Research* 4:425-432 [[Crossref](#)]
- Barda, B. D., Rinaldi, L., Ianniello, D., Zepherine, H., Salvo, F., Sadutshang, T., Cringoli, G., Clementi, M., and Albonico, M. (2013). Mini-FLOTAC, an innovative direct diagnostic technique for intestinal parasitic infections: Experience from the field. *PLoS Neglected Tropical Disease.*, 7, e2344. [[Crossref](#)]
- Burlandy-Soares, L. C., Souza-Dias, L. C., Kanamura, H. Y., Oliveira, E. J., and Ciaravolo, R. M. (2013). Schistosomiasis mansoni: follow-up of control program based on parasitologic and serologic methods in a Brazilian community of low endemicity. *Mem Inst Oswaldo Cruz*; 98:853-859. [[Crossref](#)]
- Carvalho, G. C., Marques, L. H., Gomes, L. I., Rabello, A., Ribeiro, L. C., and Scopel, K. K., (2012). Polymerase chain reaction for the evaluation of *Schistosoma mansoni* infection-endemicity endemicity areas of Minas Gerais, Brazil. *Mem Inst Oswaldo Cruz*; 107:899-902. [[Crossref](#)]
- Catherine, F. P., Samson, N. G., Kenneth, W. B., Michele, W. P., Miriam, P. R., and Lewie, Z., (2018). Biopsy and Histopathologic Diagnosis of Oral Premalignant and Malignant Lesions. 74: 3.
- Cesari, I, M., Ballen, D. E., Mendoza, L., Ferrer, A., Pointier, J. P., and Kombila M, (2018). Immunoblot analysis of membrane antigens of *Schistosoma mansoni*, *Schistosoma haematobium* against *Schistosoma*-infected patient sera. *Parasitological Research*; 106:1225-1231. [[Crossref](#)]
- Chernet, A., Kling, K.; Sydow, V., Kuenzli, E., Hatz, C., Utzinger, J., Van Lieshout, L., Marti, H., Nickel, B., and Labhardt, N. D. (2017). Accuracy of diagnostic tests for *Schistosoma mansoni* infection in asymptomatic Eritrean refugees: Serology and point-of-care circulating cathodic antigen against stool microscopy. *Clinical and Infectious Diseases*. 65, 568-574. [[Crossref](#)]
- Cioli, D., and Pica-Mattocchia, L., (2013). Praziquantel. *Parasitological Research*, 90(1):3-9. [[Crossref](#)]
- Colley, D. G., Cook, J. A., Freeman, G. L., Bartholomew, R. K., and Jordan, P. (2017). Immune responses during human Schistosomiasis mansoni. I. In vitro lymphocyte blastogenic responses to heterogeneous antigenic preparations from schistosome eggs, worms, and cercariae. *International Allergy and Applied Immunology*. 53: 420-433. [[Crossref](#)]
- Corstjens, P., de Dood, C., Kornelis, D., Fat, E., Wilson, R., Kariuki, T., Nyakundi, R. K., Loverde, P. T., Abrams, W. R., and Tanke, H. J. (2014). Tools for diagnosis, monitoring, and screening of *Schistosoma* infections utilizing lateral-flow-based assays and upconverting phosphor labels. *Parasitology*, 141, 1841-1855. [[Crossref](#)]

- Coulibaly, J. T., Goran., E. K., Utzinger, J., Doenhoff, M.J.; Dawson, E.M. (2013). A new rapid diagnostic test for detection of anti-Schistosoma mansoni and anti-Schistosoma haematobium antibodies. *Parasitology Vectors*, 6, 29-33. [\[Crossref\]](#)
- Deelder, A. M., Kornelis D, Van Marck EA, Eveleigh PC, Van Egmond JG, (2020). Schistosoma mansoni: characterization of two circulating polysaccharide antigens and the immunological response to these antigens in mouse, hamster, and human infections. *Experimental Parasitology*, 50: 16-32. [\[Crossref\]](#)
- Deelder AM. (2013). Immunology of experimental infections with Schistosoma mansoni in the Swiss mouse and Fasciola hepatica in the rabbit. *Acta Leiden*; 39:107.
- Deelder, A. M., Z. L. Qian, P. G. Kremsner, L. Acosta, A. L. T. Rabello, P. Enyong, P. P. Simarro, E. C. M. Van Etten, F. W. Krijer, J. P. Rotmans, Y. E. Fillie, N. De Jonge, A. M. Agnew, and L. Van Lieshout. (2014). Quantitative diagnosis of Schistosoma infections by measurement of circulating antigens in serum and urine. *Tropical and Geographical Medicine*. 46:233-238
- Devaney, E., Winter, A. D. and Britton, C. (2010). MicroRNAs: a role in drug resistance in parasitic nematodes. *Trends in Parasitology*, 26:428-433. [\[Crossref\]](#)
- Doenhoff, M. J., Butterworth AE, Hayes RJ, Sturrock RF, Ouma JH, Koech D, (2013). Seroepidemiology and serodiagnosis of Schistosomiasis in Kenya using crude and purified egg antigens of Schistosoma mansoni in ELISA. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 87:42-48. [\[Crossref\]](#)
- Doenhoff MJ, Butterworth AE, Hayes RJ, Sturrock RF, Ouma JH, Koech D, (2016). Seroepidemiology and serodiagnosis of schistosomiasis in Kenya using crude and purified egg antigens of Schistosoma mansoni in ELISA. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 87:42-48. [\[Crossref\]](#)
- Dunne DW, Bain J, Lillywhite J, Doenhoff MJ. (2014). The stage-, strain- and species-specificity of a Schistosoma mansoni egg antigen fraction (CEF6) with serodiagnostic potential. *Trans R Soc Trop Med Hyg*; 78:460-470 [\[Crossref\]](#)
- Dunne DW, Butterworth AE, Fulford AJ, (2017). Immunity after treatment of human Schistosomiasis: association between IgE antibodies to adult worm antigens and resistance to reinfection. *European Journal of Immunology*; 22: 1483-1494. [\[Crossref\]](#)
- El Ganayni, G. A., and M. E. Youssef. (2012). Evaluation of adult Schistosoma mansoni and cercarial antigens in serodiagnosis of Schistosomiasis using IHAT and ELISA. *Journal of Egyptian Society of Parasitology*. 22:555-560
- El Ridi, A. M., N. T. Nasr, M. M. Ismail, H. H. Hamadto, S. M. Rashed, and El Saiid, A. (2016). Comparative studies on indirect haemagglutination test and enzyme linked immunosorbent assay for diagnosis of Schistosomiasis. *Journal of Egyptian Society of Parasitology*. 16:587-592
- Enk MJ, Oliveira e Silva G, Rodrigues NB. (2012). Diagnostic accuracy and applicability of a PCR system for the detection of Schistosoma mansoni DNA in human urine samples from an endemic area. *PLoS One*; 7: e38947. [\[Crossref\]](#)
- Eriksson, C., Kokkonen H, Johansson M, Hallmans G, Wadell G, Dahlqvist SR (2011). Autoantibodies predate the onset of systemic lupus erythematosus in northern Sweden. *Arthritis Research and Therapy*., 13(1): R30. [\[Crossref\]](#)
- Erkol B, Legesse1M, Balcha1F, Medhin1G, Sori2T. Diagnosis efficacy of Indirect Hemagglutination Test in relation to kato method for diagnosis of Schistosomiasis mansoni. *Ethiopian Journal of Health Science*. 19.:2-14. [\[Crossref\]](#)
- Feldmeier, H., and D. W. Buttner. (2013). Immunodiagnosis of Schistosomiasis haematobium and Schistosomiasis mansoni in man. Application of crude extracts from adult worms and cercariae in the IHA and the ELISA. *Zentralbloti Bakteriologische Mikrobiologie Hygiene*. 255:413-421. [\[Crossref\]](#)
- Fitzsimmons, C. M., Stewart, T. J., Karl, F. H., Jane, L. G., Maria, Y., and David, W. D. (2005). Human IgE response to the Schistosoma haematobium 22·6 kDa antigen, *Parasite Immunology*, 26: 371-376. [\[Crossref\]](#)
- Friedman LS. (2004). Controversies in liver biopsy: who, where, when, how, why? *Curr Gastroenterol Rep* 6:30-36. [\[Crossref\]](#)
- Ghandour AM, Tricker K, Doenhoff MJ, al-Robai AA, Banaja AA. (2017). An enzyme-linked immunosorbent assay using Schistosoma mansoni purified egg antigen for the diagnosis of Schistosomiasis in Saudi Arabia. *Transmission*

- and Royal Society Tropical Medicinal and Hygiene. 91:287-289 [\[Crossref\]](#)
- Gray DJ, Ross AG, Li YS, McManus DP. (2021). Diagnosis and management of schistosomiasis. *Biomedical Journal.*; 342: d2651. [\[Crossref\]](#)
- Grenfell RF, Martins W, Enk M, Almeida A, Siqueira L, SilvaMoraes V, Oliveira E, Carneiro NF, Coelho PM, (2013). *Schistosoma mansoni* in a low-prevalence area in Brazil: the importance of additional methods for the diagnosis of hard-to-detect individual carriers by low-cost immunological assays. *Members of Institutional Oswaldo Cruz.* 108: 328-334. [\[Crossref\]](#)
- Gryseels B, Polman K, Clerinx J, Kestens L (2016). The distribution of *Schistosoma mansoni* in the Rusizi plain (Burundi). *Anal of Tropical Medicine and Parasitology*, 82: 581-590 [\[Crossref\]](#)
- Guegan H, Fillaux J, Charpentier E, Robert-Gangneux F, Chauvin P, Guemas E, (2019) Real-time PCR for diagnosis of imported schistosomiasis. *PLoS Neglected Tropical Diseases.* 13(9): 126-139. [\[Crossref\]](#)
- Gui, M., M. A. Idris, Y. E. Shi, A. Muhling, and A. Ruppel. (2019). Reactivity of *Schistosoma japonicum* and *S. mansoni* antigen preparations in indirect haemagglutination (IHA) with sera of patients with homologous and heterologous Schistosomiasis. *Annals of Tropical Medicine Parasitology.* 85:599-604. [\[Crossref\]](#)
- Hagan P, Blumenthal UJ, Dunn D, Simpson AJ & Wilkins HA. (2011). Human IgE, IgG4, and resistance to reinfection with *Schistosoma haematobium*. *Nature*; 349: 243- 245. [\[Crossref\]](#)
- Hamburger J, Turetski T, Kapeller I, Deresiewicz R. (2017). Highly repeated short DNA sequences in the genome of *Schistosoma mansoni* were recognized by a species-specific probe. *Molecular Biochemistry and Parasitology.*; 44: 73-80. [\[Crossref\]](#)
- Hargraves MM (2019). Production in vitro of LE cell phenomenon: use of normal bone marrow elements and blood plasma from patients with acute disseminated lupus erythematosus. *Mayo Clinical Protocol*, 24(9): 234:237.
- Hawkins, K.R.; Cantera, J.L.; Storey, H.L.; Leader, B.T.; de Los Santos, T. (2016). Diagnostic tests to support late-stage control programs for Schistosomiasis and soil-transmitted helminthiasis. *PLoS Neglected Tropical Diseases*, 1-15. [\[Crossref\]](#)
- Hervé M, Angeli V, Pinzar E, Wintjens R, Faveeuw C, Narumiya S, (2013). Pivotal roles of the parasite PGD2 synthase and of the host D prostanoid receptor 1 in schistosome immune evasion. *European Journal of Immunology*, 33 (10):2764-72. [\[Crossref\]](#)
- Hinz, R.; Schwarz, N.G.; Hahn, A.; Frickmann, H. (2017). Serological approaches for the diagnosis of schistosomiasis-A review. *Molecular and Cellular Probes*, 31, 2-21. [\[Crossref\]](#)
- Infurnari, L.; Galli, L.; Bigoloni, A.; Carbone, A.; Chiappetta, S.; Sala, A.; Ceserani, N.; Lazzarin, A.; Castagna, A.; Gaiera, G. (2017) The use of circulating cathodic antigen rapid test and serology for diagnosis of active *Schistosoma mansoni* infection in migrants in Italy, a non-endemic country: a cross-sectional study. *Members of Institutional Oswaldo Cruz*, 112, 452-455. [\[Crossref\]](#)
- John N. C, (2014): Epidemiology and transmission patterns of *Schistosoma haematobium* infections in central Ebonyi State, Nigeria. *International journal of natural and applied sciences*, 2(3): 219 - 224.
- Kanamura HY, Hoshino-Shimizu S, Camargo ME, Silva LC. (2019). Class-specific antibodies and fluorescent staining patterns in acute and chronic forms of Schistosomiasis mansoni. *American Journal of Tropical Medicine and Hygiene*; 28: 242-248 [\[Crossref\]](#)
- Karkera BV, Shivakumar BN, Mohammed A, Vidya M, Nandaprasad S, Hemanth M. (2011). Biopsy: Clinical implications. *Journal of Dental and Oral Hygiene*; 3(8): 106-108
- King, C.; Bertsch, D. (2013). Meta-analysis of urine hema dipstick diagnosis of *Schistosoma haematobium* infection, including low prevalence and previously-treated populations. *PLoS Neglected Tropical Diseases.* 7, e2431. [\[Crossref\]](#)
- Kinkel H.F., Dittrich B. Baumer and T Weitzel, (2012). Evaluation of eight serological tests for diagnosis of imported Schistosomiasis. *Clinical Vaccine and Immunology*, 19:948-953. [\[Crossref\]](#)
- Knopp S, Person B, Ame SM, (2019). Evaluation of integrated interventions layered on mass drug administration for urogenital Schistosomiasis elimination: a cluster-randomized trial. *Lancet Glob Health*; 7: e1118-29. [\[Crossref\]](#)

- Krunic AL, Wang LC, Soltani K, Weitzul S, Taylor RS. (2014). Digital anesthesia with epinephrine: An old myth revisited. *Journal of American Academic Dermatology* 51:755-759. [[Crossref](#)]
- Kumaraswamy KL, Vidhya M, Rao PK, Mukunda A. (2012). Oral biopsy: Oral pathologist's perspective. *Journal of Cancer and Research in Therapy*, 8: 192-8. [[Crossref](#)]
- Lalonde D, Bell M, Benoit P, Sparkes G, Denkler K, Chang P. A multicenter (2015) prospective study of 3,110 consecutive cases of elective epinephrine use in the fingers and hand: The Dalhousie Project clinical phase. *Journal of Hand Surgery in America*. 30:1061-7. [[Crossref](#)]
- Lindholz, C.G.; Favero, V.; Verissimo, C.d.M.; Candido, R.R.F.; de Souza, R.P.; dos Santos, R.R.; Morassutti, A.L.; Bittencourt, H.R.; Jones, M.K.; St. Pierre, T.G.; (2018). Study of diagnostic accuracy of Helmintex, Kato-Katz, and POC-CCA methods for diagnosing intestinal Schistosomiasis in Candela, a low-intensity transmission area in northeastern Brazil. *PLoS Neglected Tropical Diseases*. 12: 234-241 [[Crossref](#)]
- Lodh N, Mwansa JC, Mutengo MM, Shiff CJ. (2013). Diagnosis of *Schistosoma mansoni* without the stool: comparison of three diagnostic tests to detect *Schistosoma mansoni* infection from filtered urine in Zambia. *American Journal of Tropical Medicine and Hygiene*; 89:46-50. [[Crossref](#)]
- Luciana, I. G., Martin, J. E. and Ana R. (2013). Diagnosing schistosomiasis: where are we. *Revista da Sociedade Brasileira de Medicina Tropical*. [[Crossref](#)]
- Luciana, I. G., Martin, J. E. and Ana R. (2013). Diagnosing Schistosomiasis: where are we. *Revista da Sociedade Brasileira de Medicina Tropical*. [[Crossref](#)]
- McManus DP, Gray DJ, Li Y, Feng Z, Williams GM, Stewart D, Rey-Ladino J, Ross AG. (2010). Schistosomiasis in the People's Republic of China: the era of the Three Gorges. *Clinical Microbiology and Review*; 23(2):442-466. [[Crossref](#)]
- Meurs L, Brienen E, Mbow M (2015). Is PCR the next reference standard for the diagnosis of Schistosomiasis in stool? A comparison with microscopy in Senegal and Kenya. *PLoS of Neglected Tropical Diseases*; 9:1-16. [[Crossref](#)]
- Mutalik S. (2018). How to make local anesthesia less painful? *Journal of Cutanity Aesthetic Surgery*. 1:37-38. [[Crossref](#)]
- Nash TE. (2018). Antibody response to a polysaccharide antigen present in the schistosome gut. I. Sensitivity and specificity. *American Journal of Tropical Medicine and Hygiene*. 27: 939-943. [[Crossref](#)]
- Olliaro PL, Vaillant M, Diawara A, Coulibaly JT, Garba A, (2020). Toward Measuring Schistosoma Response to Praziquantel Treatment with Appropriate Descriptors of Egg Excretion. *PLoS Neglected Tropical Diseases*. 9: e0003821. [[Crossref](#)]
- Ortu G, O. Ndayishimiye, M. Clements, D. Kayugi and C. Campbell J. R. (2017). Countrywide reassessment of *Schistosoma mansoni* infection in Burundi a urine-circulating cathode antigen rapid test: Informing the national control program. *American Journal of Tropical Medicinal Hygiene*. 96:664-673. [[Crossref](#)]
- Patton LL, Epstein JB, Kerr AR. Adjunctive techniques for oral cancer examination and lesion diagnosis: a systematic review of the literature. *Journal American Dentist Association*; 139(7): 896-905. [[Crossref](#)]
- Pearson MS, Pickering DA, McSorley HJ, (2012). Enhanced protective efficacy of a chimeric form of the schistosomiasis vaccine antigen Sm-TSP-2. *PLoS Neglected Tropical Diseases*; 6: e1564. [[Crossref](#)]
- Pennington LF, Alouffi A, Mbanefo EC, (2017). H-IPSE is a pathogen-secreted host nucleus-infiltrating protein (Infiltrin) expressed exclusively by the *Schistosoma haematobium* egg stage. *Infection Immunity*; 85: e00301-17. [[Crossref](#)]
- Perez M, Lodha R, Nouri K. Skin biopsy techniques. In: Nouri K, Leal-Khoury S. (2013). *Techniques in dermatologic surgery*. *PLoS Neglected Tropical Diseases*. 75-80.
- Polman, K., F. F. Stelma, B. Gryseels, G. J. Van Dam, I. Talla, M. Niang, L. Van Lieshout, and A. M. Deelder. (2015). Epidemiological application of circulating antigen detection in a recent *Schistosoma mansoni* focus in northern Senegal. *American Journal of Tropical Medicine Hygiene*. 53:152-157. [[Crossref](#)]
- Rabello ALT, Rocha RS, Oliveira JPM, Katz N, Lambertucci JR (2012). Stool examination and

- rectal biopsy in the diagnosis and therapeutical evaluation of Schistosomiasis mansoni. Review Institution Medicine of Tropical Medicine 34: 601-608. [[Crossref](#)]
- Rihet P, Demeure CE, Bourgois A, Prata A & Dessein AJ. (2017). Evidence for an association between human resistance to Schistosoma mansoni and high anti-larval IgE levels. European Journal of Immunology; 21: 2679-2686. [[Crossref](#)]
- Sanjay K., Prasant MC, Kedar Saraf, Kishor Patil (2014). Oral biopsy: Techniques and their importance. American Journal of Advances in Medical Science; 2(3): 42-46
- Smith H, Doenhoff M, Aitken C, Bailey W, Ji M, Dawson E, Gilis H, Spence G, Alexander C, van Gool T, (2012). Comparison of Schistosoma mansoni soluble cercarial antigens and soluble egg antigens for serodiagnosis schistosome infections. PLoS Neglected Tropical Diseases 6: e1815 [[Crossref](#)]
- Smith JH, Christie JD. (2016). The pathobiology of Schistosoma haematobium infection in humans. Human Pathology; 17: 333-45. [[Crossref](#)]
- Smithers SR, Terry RJ. (2015). The infection of laboratory hosts with cercariae of Schistosoma mansoni and the recovery of the adult worms. Parasitology; 55:695-700. [[Crossref](#)]
- Sorgho H, Bahgat M, Poda JN, Song W, Kirsten C, Doenhoff MJ, (2015). Serodiagnosis of Schistosoma mansoni infections in an endemic area of Burkina Faso: performance of several immunological tests with different parasite antigens. Acta Tropica; 93:169-180. [[Crossref](#)]
- Stete K, Krauth SJ, Coulibaly JT, Knopp S, Hattendorf J, (2019). Dynamics of Schistosoma haematobium egg output and associated infection parameters following treatment with praziquantel in school-aged children. Parasitology Vectors.; 5: 298. [[Crossref](#)]
- Thomson CJ, Lalonde DH, Denkler KA, Feicht AJ. (2017). A critical look at the evidence for and against elective epinephrine use in the finger. Plast Reconstr Surg;119: 260-266. [[Crossref](#)]
- Towbin, H., T. Staehelin, and J. Gordon. (2019). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. Procedure of National Academic Science in United State of America. 76:4350-4354 [[Crossref](#)]
- Tsang VC, Hillyer GV, Noh J, Vivas-Gonzalez BE, Ahn LH, Pilcher JB, (2021). Geographic clustering and seroprevalence of Schistosomiasis in Puerto Rico. American Journal of Tropical Medicine and Hygiene; 56:107-112. [[Crossref](#)]
- Utzinger, J.; Becker, S.L.; van Lieshout, L.; van Dam, G.J.; Knopp, S. (2015). New diagnostic tools in Schistosomiasis. Clinical Microbiology and Infection. 21, 529-542. [[Crossref](#)]
- Van Etten, L., C. Folman, T. Eggelte, P. Kreamsner, and A. M. Deelder. (2014). Rapid diagnosis of Schistosomiasis by antigen detection in urine with a reagent strip. Journal of Clinical and Microbiology. 32:2404-2406 [[Crossref](#)]
- Van Gool T, Vetter H, Vervoort T, Doenhoff MJ, Wetsteyn J, Overboosch D. (2012). Serodiagnosis of imported Schistosomiasis by a combination of a commercial Indirect Hemagglutination Test with Schistosoma mansoni adult worm antigens and an Enzyme-Linked Immunosorbent Assay with S. mansoni eggs antigens. Journal of Clinical Microbiology. 40: 3432 - 3437. [[Crossref](#)]
- Van Lieshout, L.; Polderman, A.M.; Deelder, A.M. (2017) Immunodiagnosis of schistosomiasis by determination of the circulating antigens CAA and CCA, in particular in individuals with recent or light infections. Acta Tropica 77, 69-80. [[Crossref](#)]
- Vendrame CM, Carvalho MD, Yamamoto CR, Nakhle MC, Carvalho SA, Chieffi PP, (2021). Evaluation of anti-Schistosoma mansoni IgG antibodies in patients with chronic Schistosomiasis mansoni before and after specific treatment. Review Institutional Medicine in Tropics Sao Paulo. 43: 153-159 [[Crossref](#)]
- Webster M, Fallon PG, Fulford AJ, (2019). IgG4 and IgE responses to Schistosoma mansoni adult worms after treatment. Journal of Infectious Diseases. 175: 493-4. [[Crossref](#)]
- Wen LY, Chen JH, Ding JZ, Zhang JF, Lu SH, Yu LL, (2019). Evaluation of the applied value of the dot immunogold filtration assay (DIGFA) for rapid detection of anti-Schistosoma japonicum antibody. Acta Tropica. 96:142-147. [[Crossref](#)]
- Whitty CJ, Mabey DC, Armstrong M, Wright SG, and Chiodini PL. 2018; Presentation and outcome of 1107 cases of Schistosomiasis from Africa diagnosed in a non-endemic country. Transmission Royal Society of Tropical Medicine and Hygiene. 94: 531- [[Crossref](#)]

- Wilhelmi BJ, Blackwell SJ, Miller JH, Mancoll JS, Dardano T, Tran A, (2011). Do not use epinephrine in digital blocks: Myth or truth? *Plast Reconstr Surg.* 107:393-397 [[Crossref](#)]
- Worrell, C.M.; Bartoces, M.; Karanja, D.M.S.; Ochola, E.A.; Matete, D.O.; Mwinzi, P.N.M.; Montgomery, S.P.; Secor, W.E. (2015). Cost analysis of tests for the detection of *Schistosoma mansoni* infection in children in Western Kenya. *American Journal Tropical Medicine and Hygiene.* 92, 1233-1239. [[Crossref](#)]
- Yu JM, de Vlas SJ, Jiang QW, Gryseels B. (2017). Comparison of the Kato-Katz technique, hatching test, and indirect hemagglutination assay (IHA) for the diagnosis of *Schistosoma japonicum* infection in China. *Parasitological International;* 56:45-49. [[Crossref](#)]
- Zhou YB, Yang MX, Wang QZ, Zhao GM, Wei JG, Peng WX, (2017). Field comparison of undiagnostic and parasitological techniques for the detection of *Schistosomiasis japonica* in the People's Republic of China. *American Journal of Tropical Medicine and Hygiene.* 76:1138-1143 [[Crossref](#)]
- Zhu YC. (2015). Immunodiagnosis and its role in *Schistosomiasis* control in China: a review. *Acta Tropica;* 96:130-136 [[Crossref](#)]